arXiv:1405.5278v1 [cs.IT] 21 May 2014

The weight distribution of a family of p-ary cyclic codes

Long Yu, Hongwei Liu*

School of Mathematics and Statistics, Central China Normal University, Wuhan, Hubei 430079, China

Abstract

Let m, k be positive integers, p be an odd prime and π be a primitive element of $\mathbb{F}_{p^m}$. In this paper, we determine the weight distribution of a family of cyclic codes $\mathcal{C}_t$ over $\mathbb{F}_p$, whose duals have two zeros π^{-t} and $-\pi^{-t}$, where t satisfies $t \equiv \frac{p^k+1}{2}p^\tau \pmod{\frac{p^m-1}{2}}$ for some $\tau \in \{0, 1, \cdots, m-1\}$.

Key Words Cyclic code, quadratic form, weight distribution.

1 Introduction

Let p be an odd prime and q be a power of p. An $[n, k, d]$ linear code over the finite field $\mathbb{F}_q$ is a k-dimensional subspace of $\mathbb{F}_q^n$ with minimum Hamming distance d. A linear code $\mathcal{C}$ of length n is called cyclic if $(c_0, c_1, \cdots, c_{n-1}) \in \mathcal{C}$ implies that $(c_{n-1}, c_0, \cdots, c_{n-2}) \in \mathcal{C}$. By identifying a codeword $(c_0, c_1, \cdots, c_{n-1}) \in \mathcal{C}$ with the polynomial

$$c_0 + c_1X + \cdots + c_{n-1}X^{n-1} \in \mathbb{F}_q[X]/(X^n - 1),$$

a cyclic code $\mathcal{C}$ of length n over $\mathbb{F}_q$ corresponds to an ideal of $\mathbb{F}_q[X]/(X^n-1)$. The monic generator $g(X)$ of this ideal is called the generator polynomial of $\mathcal{C}$, which satisfies that $g(X)|(X^n-1)$. The polynomial $h(X) = (X^n - 1)/g(X)$ is referred to as the parity-check polynomial of $\mathcal{C}$ [15].

Let A_i denote the number of codewords with Hamming weight i in a linear code $\mathcal{C}$ of length n. The weight enumerator of $\mathcal{C}$ is defined by

$$A_0 + A_1X + A_2X^2 + \cdots + A_nX^n, \quad \text{where } A_0 = 1.$$

The sequence $(A_0, A_1, \cdots, A_n)$ is called the weight distribution of the code $\mathcal{C}$. In general the weight distribution of cyclic codes are difficult to be determined and they are known only for a few special classes. There are some results on the weight distribution of cyclic codes whose duals have two or more zeros(see [2, 3, 5, 6, 9, 12, 13, 16, 18–28]).

*Corresponding author.
Email address: hwliu@mail.ccnu.edu.cn (Hongwei Liu), longyu@mails.ccnu.edu.cn (Long Yu).

Let $\mathbb{Z}_m$ be the residue ring modulo integer m, Γ_i be the p-cyclotomic coset modulo p^m-1 containing i, i.e.,

$$\Gamma_i = \{i \cdot p^j \pmod{p^m-1} \mid j = 0, 1, \cdots, m-1\},$$

where $i \in \mathbb{Z}_{p^m-1}$. A subset $\{i_1, i_2, \cdots, i_r\}$ of $\{0, 1, \cdots, p^m-2\}$, where $r \geq 1$, is called a *complete set of representatives* of all p-cyclotomic cosets modulo p^m-1 if $\Gamma_{i_1}, \Gamma_{i_2}, \cdots, \Gamma_{i_r}$ are pairwise disjoin and $\bigcup_{k=1}^{r} \Gamma_{i_k} = \{0, 1, \cdots, p^m-2\}$.

Throughout this paper, we assume that m, k are positive integers, π is a primitive element of $\mathbb{F}_{p^m}$ and ζ_p is a primitive p-th unity root. Let t be an integer such that $(\pi^t)^{p^i} \neq -\pi^t$ for all $i \in \mathbb{Z}_m$. Let $h_1(x)$ and $h_2(x)$ be the minimal polynomials of π^{-t} and $-\pi^{-t}$ over $\mathbb{F}_p$, respectively. In this paper, we let $h(x) = h_1(x)h_2(x)$ and $\mathcal{C}_t$ be the cyclic code with parity-check polynomial $h(x)$. By the well-known Delsarte's Theorem [1], cyclic code $\mathcal{C}_t$ can be expressed as

$$\mathcal{C}_t = \left\{ \mathbf{c}(a,b) = \left(\mathrm{Tr}_1^m(a\pi^{ti} + b(-\pi^t)^i)\right)_{i=0}^{p^m-2} \mid a, b \in \mathbb{F}_{p^m} \right\}, \tag{1.1}$$

where $\mathrm{Tr}_1^m(\cdot)$ is the trace function from $\mathbb{F}_{p^m}$ to $\mathbb{F}_p$. In particular, in the case of $t = 1$, Vega and Wolfmann in [19] have studied cyclic code $\mathcal{C}_1$. They constructed this class of cyclic codes from the direct sum of two one-weight irreducible cyclic codes for odd m and proved that $\mathcal{C}_1$ has only two nonzero weights. In [14], Ma et al. further investigated cyclic code $\mathcal{C}_1$ for the case of even m, and gave the weight distribution of the code $\mathcal{C}_1$.

The goal of this paper is to determine the weight distribution of a family of cyclic codes $\mathcal{C}_t$ over $\mathbb{F}_p$ defined by (1.1) in the case of $t \equiv \frac{p^k+1}{2}p^\tau \pmod{\frac{p^m-1}{2}}$ for some $\tau \in \mathbb{Z}_m$. By applying the value distribution of the exponential sum $\sum_{x \in \mathbb{F}_{p^m}} \zeta_p^{\mathrm{Tr}_1^m(\alpha x^{p^k+1})}$, $\alpha \in \mathbb{F}_{p^m}$, which is given in [4, 7, 10], we obtain the value distribution of the exponential sum $\sum_{u \in \mathbb{F}_p^*} \sum_{x \in \mathbb{F}_{p^m}} (\zeta_p^{u\mathrm{Tr}_1^m((a+b)x^{p^k+1})} + \zeta_p^{u\mathrm{Tr}_1^m((a-b)x^{p^k+1})})$, $a,\ b \in \mathbb{F}_{p^m}$. Based on these results, we characterize the weight distribution of the code $\mathcal{C}_t$ defined by (1.1).

This paper is organized as follows. Section **2** gives some basic definitions and results over finite fields. In Section **3**, we determine the weight distribution of a family of cyclic codes $\mathcal{C}_t$ defined by (1.1) over $\mathbb{F}_p$.

2 Preliminaries

Let $\mathbb{F}_q$ be a finite field and $\mathbb{F}_q^* = \mathbb{F}_q \setminus \{0\}$, where q is a power of an odd prime p. In the following, we give a brief introduction to the theory of quadratic forms over finite fields, which is needed to calculate the weight distribution of the cyclic codes in the next section. Quadratic forms have been well studied ([10]), and they have many applications in sequence design and coding theory.

Definition 2.1. *Let $\{\omega_1, \omega_2, \cdots, \omega_s\}$ be a basis for $\mathbb{F}_{q^s}$ over $\mathbb{F}_q$ and $x = \sum_{i=1}^{s} x_i\omega_i$, where $x_i \in \mathbb{F}_q$. A function $f(x)$ from $\mathbb{F}_{q^s}$ to $\mathbb{F}_q$ is called quadratic form if it can be represented as*

$$f(x) = f(\sum_{i=1}^{s} x_i\omega_i) = \sum_{1 \leq i \leq j \leq s} a_{ij}x_ix_j, \qquad a_{ij} \in \mathbb{F}_q.$$

The rank of the quadratic form $f(x)$ is defined as the codimension of the $\mathbb{F}_q$-vector space

$$V = \{x \in \mathbb{F}_{q^s} \mid f(x+z) - f(x) - f(z) = 0, \quad \text{for all } z \in \mathbb{F}_{q^s}\},$$

denotes by rank (f). Then $|V| = q^{s-\mathrm{rank}(f)}$.

For a quadratic form $f(x)$ with s variables over $\mathbb{F}_q$, there exists a symmetric matrix A of order s over $\mathbb{F}_q$ such that $f(x) = XAX'$, where $X = (x_1, x_2, \cdots, x_s) \in \mathbb{F}_q^s$ and X' denotes the transpose of X. It is known that there exists a nonsingular matrix T over $\mathbb{F}_q$ such that TAT' is a diagonal matrix [10]. Making a nonsingular linear substitution $X = ZT$ with $Z = (z_1, z_2, \cdots, z_s) \in \mathbb{F}_q^s$, we have

$$f(x) = Z(TAT')Z' = \sum_{i=1}^{r} a_i z_i^2, \qquad a_i \in \mathbb{F}_q^*,$$

where r is the rank of $f(x)$.

Let $v_2(j)$ denote the 2-adic valuation of integer j (i.e., the maximal power of 2 dividing j). We denote the quadratic character of $\mathbb{F}_{p^m}$ by

$$\eta_m(x) = \begin{cases} 0, & \text{if } x = 0; \\ x^{\frac{p^m-1}{2}}, & \text{if } x \in \mathbb{F}_{p^m}^*. \end{cases}$$

Lemma 2.2. [4, 7] *Let k, m be positive integers. Then*

$$\gcd(p^k+1, p^m-1) = \begin{cases} p^{\gcd(k,m)}+1, & \text{if } v_2(m) > v_2(k); \\ 2, & \text{otherwise.} \end{cases}$$

The following lemma could be found in [4, 7, 10].

Lemma 2.3. [4, 7, 10] *Let $T_\alpha(x) = \sum\limits_{x \in \mathbb{F}_{p^m}} \zeta_p^{\mathrm{Tr}_1^m(\alpha x^{p^k+1})}$, where k is a positive integer. Then for any $\alpha \in \mathbb{F}_{p^m}$,*

- *if $v_2(k) \geq v_2(m)$, then*

$$T_\alpha(x) = \begin{cases} \eta_m(\alpha)(-1)^{m-1} p^{\frac{m}{2}}, & \text{if } p \equiv 1 \pmod 4; \\ \eta_m(\alpha)(\sqrt{-1})^m (-1)^{m-1} p^{\frac{m}{2}}, & \text{if } p \equiv 3 \pmod 4. \end{cases}$$

- *if $v_2(k)+1 = v_2(m)$, then*

$$T_\alpha(x) = \begin{cases} p^{\frac{m}{2}+d}, & \frac{p^m-1}{p^d+1} \text{ times;} \\ -p^{\frac{m}{2}}, & \frac{p^d(p^m-1)}{p^d+1} \text{ times;} \\ p^m, & 1 \text{ time.} \end{cases}$$

- *if $v_2(k)+1 < v_2(m)$, then*

$$T_\alpha(x) = \begin{cases} -p^{\frac{m}{2}+d}, & \frac{p^m-1}{p^d+1} \text{ times;} \\ p^{\frac{m}{2}}, & \frac{p^d(p^m-1)}{p^d+1} \text{ times;} \\ p^m, & 1 \text{ time.} \end{cases}$$

3 The weight distribution of the code $\mathcal{C}_t$

In this section, we let $n = p^m - 1$. For a given positive divisor l of m, the trace function from $\mathbb{F}_{p^m}$ to $\mathbb{F}_{p^l}$ is defined by $\mathrm{Tr}_l^m(x) = \sum_{i=0}^{\frac{m}{l}-1} x^{p^{li}}$, where $x \in \mathbb{F}_{p^m}$. Let SQ denote the set of square elements in $\mathbb{F}_{p^m}^*$, SQ_p denote the set of square elements in $\mathbb{F}_p^*$ and let u_p be a primitive element in $\mathbb{F}_p$. In the following, we compute the weight of the codeword $\mathbf{c}(a,b) \in \mathcal{C}_t$ defined by (1.1).

$$\begin{aligned}
wt(\mathbf{c}(a,b)) &= \#\{0 \le i \le p^m - 2 : c_i \neq 0\} \\
&= n - \frac{1}{p} \sum_{i=0}^{p^m-2} \sum_{u \in \mathbb{F}_p} \zeta_p^{u\mathrm{Tr}_1^m(a\pi^{ti} + b(-\pi^t)^i)} \\
&= n - \frac{1}{p} \sum_{u \in \mathbb{F}_p} \sum_{i=0}^{\frac{p^m-3}{2}} \left(\zeta_p^{u\mathrm{Tr}_1^m(a\pi^{2ti} + b(-\pi^t)^{2i})} + \zeta_p^{u\mathrm{Tr}_1^m(a\pi^{(2i+1)t} + b(-\pi^t)^{(2i+1)})} \right) \\
&= n - \frac{1}{p} \sum_{u \in \mathbb{F}_p} \sum_{x \in SQ} \left(\zeta_p^{u\mathrm{Tr}_1^m(ax^t + bx^t)} + \zeta_p^{u\mathrm{Tr}_1^m(a(\pi x)^t - b(\pi x)^t)} \right) \\
&= n - \frac{1}{2p} \sum_{u \in \mathbb{F}_p} \sum_{x \in \mathbb{F}_{p^m}^*} \left(\zeta_p^{u\mathrm{Tr}_1^m(ax^{2t} + bx^{2t})} + \zeta_p^{u\mathrm{Tr}_1^m(a\pi^t x^{2t} - b\pi^t x^{2t})} \right) \\
&= p^m - p^{m-1} - \frac{1}{2p} \sum_{u \in \mathbb{F}_p^*} \sum_{x \in \mathbb{F}_{p^m}} \left(\zeta_p^{u\mathrm{Tr}_1^m((a+b)x^{2t})} + \zeta_p^{u\mathrm{Tr}_1^m((a-b)\pi^t x^{2t})} \right). \qquad (3.1)
\end{aligned}$$

Therefore, we have the following lemma.

Lemma 3.1. *With the notations given above. If $t, e \in \mathbb{Z}_{p^m-1}$ are two positive integers such that $t \equiv ep^\tau \pmod{\frac{p^m-1}{2}}$ for some $\tau \in \mathbb{Z}_m$, then the cyclic codes $\mathcal{C}_t$ and $\mathcal{C}_e$ defined by (1.1) have the same weight distribution.*

Proof. By (3.1), we have that the weight distribution of $\mathcal{C}_t$ and $\mathcal{C}_e$ are respectively determined by the value distribution of

$$\Delta_t = \sum_{u \in \mathbb{F}_p^*} \sum_{x \in \mathbb{F}_{p^m}} \left(\zeta_p^{u\mathrm{Tr}_1^m((a+b)x^{2t})} + \zeta_p^{u\mathrm{Tr}_1^m((a-b)\pi^t x^{2t})} \right)$$

and

$$\Delta_e = \sum_{u \in \mathbb{F}_p^*} \sum_{x \in \mathbb{F}_{p^m}} \left(\zeta_p^{u\mathrm{Tr}_1^m((a+b)x^{2e})} + \zeta_p^{u\mathrm{Tr}_1^m((a-b)\pi^e x^{2e})} \right).$$

Let $e \equiv rp^{m-\tau} \pmod{p^m-1}$, then the integers t and r satisfy $t \equiv r \pmod{\frac{p^m-1}{2}}$, i.e. $t = r + l\frac{p^m-1}{2}$ for some integer l. Hence

$$\begin{aligned}
\Delta_t &= \sum_{u \in \mathbb{F}_p^*} \sum_{x \in \mathbb{F}_{p^m}} \left(\zeta_p^{u\mathrm{Tr}_1^m((a+b)x^{2t})} + \zeta_p^{u\mathrm{Tr}_1^m((a-b)\pi^t x^{2t})} \right) \\
&= \sum_{u \in \mathbb{F}_p^*} \sum_{x \in \mathbb{F}_{p^m}} \left(\zeta_p^{u\mathrm{Tr}_1^m((a+b)x^{2r+l(p^m-1)})} + \zeta_p^{u\mathrm{Tr}_1^m((a-b)\pi^{r+l\frac{p^m-1}{2}} x^{2r+l(p^m-1)})} \right)
\end{aligned}$$

$$
\begin{aligned}
&= \sum_{u\in\mathbb{F}_p^*}\sum_{x\in\mathbb{F}_{p^m}}\left(\zeta_p^{u\mathrm{Tr}_1^m((a+b)x^{2r})}+\zeta_p^{u\pi^{l\frac{p^m-1}{2}}\mathrm{Tr}_1^m((a-b)\pi^r x^{2r})}\right)\\
&= \sum_{u\in\mathbb{F}_p^*}\sum_{x\in\mathbb{F}_{p^m}}\left(\zeta_p^{u\mathrm{Tr}_1^m((a+b)x^{2r})}+\zeta_p^{u\mathrm{Tr}_1^m((a-b)\pi^r x^{2r})}\right)\\
&= \Delta_r
\end{aligned}
$$

where the fourth identity is obtained by $u\pi^{l\frac{p^m-1}{2}}=\pm u$. As we know, $\{\Delta_r \mid a,b\in\mathbb{F}_{p^m}\}=\{\Delta_e \mid a,b\in\mathbb{F}_{p^m}\}$, since $e\equiv rp^{m-\tau} \pmod{p^m-1}$. Therefore, the multi-sets $\{\Delta_t \mid a,b\in\mathbb{F}_{p^m}\}$ and $\{\Delta_e \mid a,b\in\mathbb{F}_{p^m}\}$ have the same value distribution. □

In this section, we study the weight distribution of cyclic codes $\mathcal{C}_t$ in the case of $t\equiv\frac{p^k+1}{2}p^\tau \pmod{\frac{p^m-1}{2}}$ for some $\tau\in\mathbb{Z}_m$. By Lemma 3.1, we have that the codes $\mathcal{C}_t$ and $\mathcal{C}_{\frac{p^k+1}{2}}$ have the same weight distribution. In order to determine the weight distribution of the code $\mathcal{C}_t$, we just need to obtain the weight distribution of the code $\mathcal{C}_{\frac{p^k+1}{2}}$. In the following, we calculate the weight distribution of cyclic code $\mathcal{C}_{\frac{p^k+1}{2}}$. By (1.1), we have

$$
\mathcal{C}_{\frac{p^k+1}{2}}=\left\{\mathbf{c}(a,b)=\left(\mathrm{Tr}_1^m(a\pi^{i\frac{p^k+1}{2}}+b(-\pi^{\frac{p^k+1}{2}})^i)\right)_{i=0}^{p^m-2} \mid a,b\in\mathbb{F}_{p^m}\right\}, \tag{3.2}
$$

whose dual has two zeros $\pi^{-\frac{p^k+1}{2}}$ and $-\pi^{\frac{p^k+1}{2}}$, where k satisfies $\pi^{\frac{p^k+1}{2}p^i}\neq-\pi^{\frac{p^k+1}{2}}$ for all $i\in\mathbb{Z}_m$. Let $h_1(x)$ and $h_2(x)$ be the minimal polynomials of $\pi^{-\frac{p^k+1}{2}}$ and $-\pi^{-\frac{p^k+1}{2}}$ over $\mathbb{F}_p$, respectively. Then we have the following lemma.

Lemma 3.2. *With the notations given above. The degree of $h_1(x)$ and $h_2(x)$ are both m.*

Proof. In order to calculate the degree of $h_1(x)$, we need to compute the size of $\Gamma_{\frac{p^k+1}{2}}$. Let the size of $\Gamma_{\frac{p^k+1}{2}}$ be l. Note that $\frac{p^k+1}{2}p^m\equiv\frac{p^k+1}{2} \pmod{p^m-1}$, then we get that $l\mid m$. On the other hand, we have

$$
\frac{p^k+1}{2}p^l\equiv\frac{p^k+1}{2} \pmod{p^m-1},
$$

which implies that

$$
2(p^m-1)\mid(p^k+1)(p^l-1). \tag{3.3}
$$

Case I, when $v_2(k)\geq v_2(m)$: By Lemma 2.2, we have $\gcd(p^k+1,p^m-1)=2$. From (3.3), we get $m\mid l$. Since $l\mid m$, hence, $m=l$.

Case II, when $v_2(k)<v_2(m)$: In this case, we obtain that $\gcd(p^k+1,p^m-1)=p^d+1$ by Lemma 2.2. From (3.3), we have $2\frac{p^m-1}{p^d+1}\mid(p^l-1)$. Since $v_2(k)<v_2(m)$, then $p^{2d}-1\mid p^m-1$, which implies that $p^d-1\mid 2\frac{p^m-1}{p^d+1}$. This shows that $p^d-1\mid p^l-1$, i.e., $d\mid l$. Let $l=hd$, $m=sd$, where $h\mid s$ since $l\mid m$. From (3.3), we have

$$
2(p^{sd}-1)\mid(p^d+1)(p^{hd}-1),
$$

which implies that $h=s$. Then $l=m$.

Similarly, we also get that the degree of $h_2(x)$ is m. □

From (3.1), the weight of $\mathbf{c}(a,b) \in \mathcal{C}_{\frac{p^k+1}{2}}$ can be expressed as

$$wt(\mathbf{c}(a,b)) = p^m - p^{m-1} - \frac{1}{2p} \sum_{u\in\mathbb{F}_p^*} \sum_{x\in\mathbb{F}_{p^m}} \left(\zeta_p^{u\mathrm{Tr}_1^m((a+b)x^{p^k+1})} + \zeta_p^{u\mathrm{Tr}_1^m((a-b)\pi^{\frac{p^k+1}{2}}x^{p^k+1})} \right). \quad (3.4)$$

3.1 The weight distribution of $\mathcal{C}_{\frac{p^k+1}{2}}$ for $v_2(m) > v_2(k)$

In this subsection, we always assume that $v_2(m) > v_2(k)$, i.e., $s = \frac{m}{d}$ is even, where $d = \gcd(m,k)$. Following the above notations, we let

$$T(a,b) = \sum_{u\in\mathbb{F}_p^*} \sum_{x\in\mathbb{F}_{p^m}} \left(\zeta_p^{u\mathrm{Tr}_1^m((a+b)x^{p^k+1})} + \zeta_p^{u\mathrm{Tr}_1^m((a-b)\pi^{\frac{p^k+1}{2}}x^{p^k+1})} \right). \quad (3.5)$$

From (3.4), the weight distribution of cyclic code $\mathcal{C}_{\frac{p^k+1}{2}}$ is completely determined by the value distribution of $T(a,b)$. To calculate the value distribution of $T(a,b)$, we need a series of lemmas. Before introducing them, we define

$$R_\alpha(x) = \sum_{u\in\mathbb{F}_p^*} \sum_{x\in\mathbb{F}_{p^m}} \zeta_p^{u\mathrm{Tr}_1^m(\alpha x^{p^k+1})},\ \alpha \in \mathbb{F}_{p^m}.$$

Lemma 3.3. *With the notations given above, we have*

$$R_\alpha(x) = (p-1) \sum_{x\in\mathbb{F}_{p^m}} \zeta_p^{\mathrm{Tr}_1^m(\alpha x^{p^k+1})}. \quad (3.6)$$

Proof. Note that $v_2(m) > v_2(k)$, then m is even and $\frac{k}{d}$ is odd. Hence, $u_p = \pi^{\frac{p^m-1}{p-1}} = \pi^{p^{m-1}+\cdots+1}$ is a square element in $\mathbb{F}_{p^m}$, i.e., $u_p \in SQ$. Since $\frac{k}{d}$ is odd and

$$u_p^{\frac{p^k+1}{2}} = u_p^{\frac{p^k-1}{2}+1} = u_p u_p^{\frac{p^d-1}{2}(p^{(\frac{k}{d}-1)d}+\cdots+1)} = u_p (u_p^{\frac{p-1}{2}})^{(p^{d-1}+\cdots+1)(p^{(\frac{k}{d}-1)d}+\cdots+1)},$$

we have that $u_p^{\frac{p^k+1}{2}} = -u_p$ for odd d, and $u_p^{\frac{p^k+1}{2}} = u_p$ for even d, i.e., $u_p^{\frac{p^k+1}{2}} = (-1)^d u_p$. Using $u_p \in SQ$, we have

$$\begin{aligned}
R_\alpha(x) &= \sum_{u\in\mathbb{F}_p^*} \sum_{x\in\mathbb{F}_{p^m}} \zeta_p^{u\mathrm{Tr}_1^m(\alpha x^{p^k+1})} \\
&= \sum_{u\in SQ_p} \sum_{x\in\mathbb{F}_{p^m}} \left(\zeta_p^{\mathrm{Tr}_1^m(\alpha(u^{\frac{1}{2}}x)^{p^k+1})} + \zeta_p^{u_p\mathrm{Tr}_1^m(\alpha(u^{\frac{1}{2}}x)^{p^k+1})} \right) \\
&= \frac{p-1}{2} \sum_{x\in\mathbb{F}_{p^m}} \left(\zeta_p^{\mathrm{Tr}_1^m(\alpha x^{p^k+1})} + \zeta_p^{u_p\mathrm{Tr}_1^m(\alpha x^{p^k+1})} \right) \\
&= \frac{p-1}{2} \sum_{x\in\mathbb{F}_{p^m}} \left(\zeta_p^{\mathrm{Tr}_1^m(\alpha x^{p^k+1})} + \zeta_p^{\mathrm{Tr}_1^m((-1)^d\alpha(u_p^{\frac{1}{2}}x)^{p^k+1})} \right)
\end{aligned}$$

$$
\begin{aligned}
&= \frac{p-1}{2}\sum_{x\in\mathbb{F}_{p^m}}\left(\zeta_p^{\mathrm{Tr}_1^m(\alpha x^{p^k+1})}+\zeta_p^{\mathrm{Tr}_1^m((-1)^d\alpha x^{p^k+1})}\right)\\
&= \begin{cases} (p-1)\sum\limits_{x\in\mathbb{F}_{p^m}}\zeta_p^{\mathrm{Tr}_1^m(\alpha x^{p^k+1})}, & \text{if } d \text{ is even;}\\ \frac{p-1}{2}\sum\limits_{x\in\mathbb{F}_{p^m}}\left(\zeta_p^{\mathrm{Tr}_1^m(\alpha x^{p^k+1})}+\zeta_p^{\mathrm{Tr}_1^m(-\alpha x^{p^k+1})}\right), & \text{if } d \text{ is odd.}\end{cases}
\end{aligned}
\tag{3.7}
$$

Since $v_2(m) > v_2(k)$, by Lemma 2.2, we have

$$\gcd(p^k+1, p^m-1) = p^d+1.$$

Note that $m = sd$ is even, then $2(p^d+1) \mid p^m-1$. Hence, we have

$$(\pi^{\frac{p^m-1}{2(p^d+1)}})^{p^k+1} = (\pi^{\frac{p^m-1}{2}})^{\frac{p^k+1}{p^d+1}} = (-1)^{\frac{p^k+1}{p^d+1}}.$$

Since $\frac{k}{d}$ is odd, then $\frac{p^k+1}{p^d+1} = p^{(\frac{k}{d}-1)d} - p^{(\frac{k}{d}-2)d} + p^{(\frac{k}{d}-3)d} - \cdots - p + 1$ is odd, which implies that $(\pi^{\frac{p^m-1}{2(p^d+1)}})^{p^k+1} = -1$. Therefore, we have

$$\sum_{x\in\mathbb{F}_{p^m}}\zeta_p^{\mathrm{Tr}_1^m(\alpha x^{p^k+1})} = \sum_{x\in\mathbb{F}_{p^m}}\zeta_p^{\mathrm{Tr}_1^m(\alpha(\pi^{\frac{p^m-1}{2(p^d+1)}}x)^{p^k+1})} = \sum_{x\in\mathbb{F}_{p^m}}\zeta_p^{\mathrm{Tr}_1^m(-\alpha x^{p^k+1})}.$$

By (3.7), we obtain

$$R_\alpha(x) = (p-1)\sum_{x\in\mathbb{F}_{p^m}}\zeta_p^{\mathrm{Tr}_1^m(\alpha x^{p^k+1})}.$$

□

Applying Lemmas 2.3 and 3.3, we have the following result.

Lemma 3.4. *With the notations given above.*

- *If $v_2(k)+1 = v_2(m)$, then*

$$R_\alpha(x) = \begin{cases} (p-1)p^m, & 1 \text{ time;}\\ -(p-1)p^{\frac{m}{2}}, & \frac{p^d(p^m-1)}{p^d+1} \text{ times;}\\ (p-1)p^{\frac{m}{2}+d}, & \frac{p^m-1}{p^d+1} \text{ times.}\end{cases}$$

- *If $v_2(k)+1 < v_2(m)$, then*

$$R_\alpha(x) = \begin{cases} (p-1)p^m, & 1 \text{ time;}\\ (p-1)p^{\frac{m}{2}}, & \frac{p^d(p^m-1)}{p^d+1} \text{ times;}\\ -(p-1)p^{\frac{m}{2}+d}, & \frac{p^m-1}{p^d+1} \text{ times.}\end{cases}$$

Lemma 3.5. *With the notations given above. Suppose (a,b) runs through $\mathbb{F}_{p^m}^2$,*

- *if* $v_2(k)+1=v_2(m)$, *then* $T(a,b)$ *takes on only the values from set of*
$\{2(p-1)p^m, (p-1)(p^m+p^{\frac{m}{2}+d}), (p-1)(p^m-p^{\frac{m}{2}}), (p-1)(p^{\frac{m}{2}+d}-p^{\frac{m}{2}}), 2(p-1)p^{\frac{m}{2}+d}, -2(p-1)p^{\frac{m}{2}}\}$.

- *if* $v_2(k)+1<v_2(m)$, *then* $T(a,b)$ *takes on only the values from set of*
$\{2(p-1)p^m, (p-1)(p^m-p^{\frac{m}{2}+d}), (p-1)(p^m+p^{\frac{m}{2}}), (p-1)(p^{\frac{m}{2}}-p^{\frac{m}{2}+d}), -2(p-1)p^{\frac{m}{2}+d}, 2(p-1)p^{\frac{m}{2}}\}$.

Proof. By (3.5) and (3.6), we have

$$T(a,b)=R_{a+b}(x)+R_{(a-b)\pi^{\frac{p^k+1}{2}}}(x).$$

We first discuss the value of $T(a,b)$ in the case of $v_2(k)+1=v_2(m)$.

Case I, when $a=b=0$: It is easy to check that $T(a,b)=2(p-1)p^m$.

Case II, when $a=b\neq 0$ or $a=-b\neq 0$: We first discuss the case of $a=b\neq 0$, since the other case can be discussed by a similar way. By Lemma 3.4, we have $R_{a+b}(x)\in\{(p-1)p^{\frac{m}{2}+d}, -(p-1)p^{\frac{m}{2}}\}$, and $R_{(a-b)\pi^{\frac{p^k+1}{2}}}(x)=(p-1)p^m$. Hence, $T(a,b)\in\{(p-1)(p^m+p^{\frac{m}{2}+d}), (p-1)(p^m-p^{\frac{m}{2}})\}$. Similarly, we get $T(a,b)\in\{(p-1)(p^m+p^{\frac{m}{2}+d}), (p-1)(p^m-p^{\frac{m}{2}})\}$ in the case of $a=-b\neq 0$.

Case III, when $a\neq b$, $a\neq -b$: By Lemma 3.4, we have $R_{a+b}(x)\in\{(p-1)p^{\frac{m}{2}+d}, -(p-1)p^{\frac{m}{2}}\}$ and $R_{(a-b)\pi^{\frac{p^k+1}{2}}}(x)\in\{(p-1)p^{\frac{m}{2}+d}, -(p-1)p^{\frac{m}{2}}\}$. Therefore, $T(a,b)\in\{2(p-1)p^{\frac{m}{2}+d}, -2(p-1)p^{\frac{m}{2}}, (p-1)(p^{\frac{m}{2}+d}-p^{\frac{m}{2}})\}$.

The case of $v_2(k)+1<v_2(m)$ can be discussed by a similar way as the case of $v_2(k)+1=v_2(m)$. This completes the proof. □

With above preparation we can determine the value distribution of the exponential sum $T(a,b)$ defined by (3.5).

Theorem 3.6. *With the notations given above. Suppose* (a,b) *runs through* $\mathbb{F}_{p^m}^2$,

- *if* $v_2(k)+1=v_2(m)$, *then the value distribution of* $T(a,b)$ *is given as follows.*

$$\begin{cases} 2(p-1)p^m, & 1 \text{ time;} \\ (p-1)(p^m+p^{\frac{m}{2}+d}), & 2\frac{p^m-1}{p^d+1} \text{ times;} \\ (p-1)(p^m-p^{\frac{m}{2}}), & 2\frac{p^d(p^m-1)}{p^d+1} \text{ times;} \\ (p-1)(p^{\frac{m}{2}+d}-p^{\frac{m}{2}}), & 2p^d(\frac{p^m-1}{p^d+1})^2 \text{times;} \\ 2(p-1)p^{\frac{m}{2}+d}, & (\frac{p^m-1}{p^d+1})^2 \text{times;} \\ -2(p-1)p^{\frac{m}{2}}, & p^{2d}(\frac{p^m-1}{p^d+1})^2 \text{times.} \end{cases}$$

- *if* $v_2(k)+1<v_2(m)$, *then the value distribution of* $T(a,b)$ *is given as follows.*

$$\begin{cases} 2(p-1)p^m, & 1 \text{ time;} \\ (p-1)(p^m-p^{\frac{m}{2}+d}), & 2\frac{p^m-1}{p^d+1} \text{ times;} \\ (p-1)(p^m+p^{\frac{m}{2}}), & 2\frac{p^d(p^m-1)}{p^d+1} \text{ times;} \\ (p-1)(p^{\frac{m}{2}}-p^{\frac{m}{2}+d}), & 2p^d(\frac{p^m-1}{p^d+1})^2 \text{times;} \\ -2(p-1)p^{\frac{m}{2}+d}, & (\frac{p^m-1}{p^d+1})^2 \text{times;} \\ 2(p-1)p^{\frac{m}{2}}, & p^{2d}(\frac{p^m-1}{p^d+1})^2 \text{times.} \end{cases}$$

Proof. We only discuss the case of $v_2(k)+1=v_2(m)$, since the other case can be discussed by a similar way. To determine the value distribution of $T(a,b)$, we define

$$N_1=\#\{a,b\in\mathbb{F}_{p^m}\mid T(a,b)=(p-1)(p^m+p^{\frac{m}{2}+d})\}, N_2=\#\{a,b\in\mathbb{F}_{p^m}\mid T(a,b)=(p-1)(p^m-p^{\frac{m}{2}})\},$$

$$N_3=\#\{a,b\in\mathbb{F}_{p^m}\mid T(a,b)=(p-1)(p^{\frac{m}{2}+d}-p^{\frac{m}{2}})\}, N_4=\#\{a,b\in\mathbb{F}_{p^m}\mid T(a,b)=2(p-1)p^{\frac{m}{2}+d}\},$$

$$N_5=\#\{a,b\in\mathbb{F}_{p^m}\mid T(a,b)=-2(p-1)p^{\frac{m}{2}}\}.$$

It is easy to check that the value $2(p-1)p^m$ happens only once. Note that the value $(p-1)(p^m+p^{\frac{m}{2}+d})$ occurs only if $R_{a+b}(x)=(p-1)p^m$, $R_{(a-b)\pi^{\frac{p^k+1}{2}}}(x)=(p-1)p^{\frac{m}{2}+d}$ or $R_{a+b}(x)=(p-1)p^{\frac{m}{2}+d}$, $R_{(a-b)\pi^{\frac{p^k+1}{2}}}(x)=(p-1)p^m$. By Lemma 3.4, if $R_{a+b}(x)=(p-1)p^m$, $R_{(a+b)\pi^{\frac{p^k+1}{2}}}(x)=(p-1)p^{\frac{m}{2}+d}$, then we have that the number of $(a,b)\in\mathbb{F}_{p^m}^2$ is $\frac{p^m-1}{p^d+1}$. By Lemma 3.4, if $R_{a+b}(x)=(p-1)p^{\frac{m}{2}+d}$, $R_{(a-b)\pi^{\frac{p^k+1}{2}}}(x)=(p-1)p^m$, we get that the number of $(a,b)\in\mathbb{F}_{p^m}^2$ is $\frac{p^m-1}{p^d+1}$. Therefore, we have $N_1=2\frac{p^m-1}{p^d+1}$. Similarly, we obtain that $N_2=2\frac{p^d(p^m-1)}{p^d+1}$. On the other hand, we have

$$\begin{aligned}
N_3 &= \#\{a,b\in\mathbb{F}_{p^m}\mid T(a,b)=R_{a+b}(x)+R_{(a-b)\pi^{\frac{p^k+1}{2}}}(x)=(p-1)(p^{\frac{m}{2}+d}-p^{\frac{m}{2}})\}\\
&= \#\{u,v\in\mathbb{F}_{p^m}\mid R_u(x)+R_v(x)=(p-1)(p^{\frac{m}{2}+d}-p^{\frac{m}{2}})\}\\
&= \#\{u,v\in\mathbb{F}_{p^m}\mid R_u(x)=(p-1)p^{\frac{m}{2}+d}, R_v(x)=-(p-1)p^{\frac{m}{2}}\}\\
&\quad +\#\{u,v\in\mathbb{F}_{p^m}\mid R_v(x)=(p-1)p^{\frac{m}{2}+d}, R_u(x)=-(p-1)p^{\frac{m}{2}}\}.
\end{aligned}$$

By Lemma 3.4, we obtain $N_3=2p^d(\frac{p^m-1}{p^d+1})^2$. Similarly, we get

$$N_4=(\frac{p^m-1}{p^d+1})^2,\quad N_5=p^{2d}(\frac{p^m-1}{p^d+1})^2.$$

This finishes the proof. □

Table 1: for the case of $v_2(k)+1=v_2(m)$

Weight	Frequency
0	1
$\frac{p-1}{2}(p^{m-1}-p^{\frac{m}{2}+d-1})$	$2\frac{p^m-1}{p^d+1}$
$\frac{p-1}{2}(p^{m-1}+p^{\frac{m}{2}-1})$	$2\frac{p^d(p^m-1)}{p^d+1}$
$\frac{p-1}{2}(2p^{m-1}-p^{\frac{m}{2}+d-1}+p^{\frac{m}{2}-1})$	$2p^d(\frac{p^m-1}{p^d+1})^2$
$(p-1)(p^{m-1}-p^{\frac{m}{2}+d-1})$	$(\frac{p^m-1}{p^d+1})^2$
$(p-1)(p^{m-1}+p^{\frac{m}{2}-1})$	$p^{2d}(\frac{p^m-1}{p^d+1})^2$

Theorem 3.7. *With the notations given above.*

- *If $v_2(k)+1=v_2(m)$, then $\mathcal{C}_{\frac{p^k+1}{2}}$ is a cyclic code over $\mathbb{F}_p$ with parameters $[p^m-1, 2m, \frac{p-1}{2}(p^{m-1}-p^{\frac{m}{2}+d-1})]$ and the weight distribution of $\mathcal{C}_{\frac{p^k+1}{2}}$ is given in Table 1.*

- *If* $v_2(k)+1 < v_2(m)$, *then* $\mathcal{C}_{\frac{p^k+1}{2}}$ *is a cyclic code over* $\mathbb{F}_p$ *with parameters* $[p^m-1, 2m, \frac{p-1}{2}(p^{m-1}-p^{\frac{m}{2}-1})]$ *and the weight distribution of* $\mathcal{C}_{\frac{p^k+1}{2}}$ *is given in Table* 2.

Table 2: for the case of $v_2(k)+1 < v_2(m)$

Weight	Frequency
0	1
$\frac{p-1}{2}(p^{m-1}+p^{\frac{m}{2}+d-1})$	$2\frac{p^m-1}{p^d+1}$
$\frac{p-1}{2}(p^{m-1}-p^{\frac{m}{2}-1})$	$2\frac{p^d(p^m-1)}{p^d+1}$
$\frac{p-1}{2}(2p^{m-1}+p^{\frac{m}{2}+d-1}-p^{\frac{m}{2}-1})$	$2p^d(\frac{p^m-1}{p^d+1})^2$
$(p-1)(p^{m-1}+p^{\frac{m}{2}+d-1})$	$(\frac{p^m-1}{p^d+1})^2$
$(p-1)(p^{m-1}-p^{\frac{m}{2}-1})$	$p^{2d}(\frac{p^m-1}{p^d+1})^2$

Proof. Combining Theorem 3.6, Lemma 3.2 and (3.4), we finish the proof. □

3.2 The weight distribution of $\mathcal{C}_{\frac{p^k+1}{2}}$ for $v_2(m) \le v_2(k)$

In this subsection, we always assume that $v_2(m) \le v_2(k)$, i.e., $s = \frac{m}{d}$ is odd, where $d = \gcd(m,k)$.

Lemma 3.8. *With the notations given above, the codes* $\mathcal{C}_1$ *and* $\mathcal{C}_{\frac{p^k+1}{2}}$ *have the same weight distribution.*

Proof. By (3.1), we have that the weight distribution of $\mathcal{C}_1$ and $\mathcal{C}_{\frac{p^k+1}{2}}$ are respectively determined by the value distribution of

$$\Delta_1 = \sum_{u\in\mathbb{F}_p^*}\sum_{x\in\mathbb{F}_{p^m}} \left(\zeta_p^{u\mathrm{Tr}_1^m((a+b)x^2)} + \zeta_p^{u\mathrm{Tr}_1^m((a-b)\pi x^2)}\right)$$

and

$$\Delta_{\frac{p^k+1}{2}} = \sum_{u\in\mathbb{F}_p^*}\sum_{x\in\mathbb{F}_{p^m}} \left(\zeta_p^{u\mathrm{Tr}_1^m((a+b)x^{p^k+1})} + \zeta_p^{u\mathrm{Tr}_1^m((a-b)\pi^{\frac{p^k+1}{2}}x^{p^k+1})}\right).$$

Since $v_2(k) \ge v_2(m)$, by Lemma 2.2, we have $\gcd(p^m-1, p^k+1) = 2$, which implies that $\{x^{p^k+1} \mid x\in\mathbb{F}_{p^m}\} = \{x^2 \mid x\in\mathbb{F}_{p^m}\}$. Hence

$$\begin{aligned}\Delta_{\frac{p^k+1}{2}} &= \sum_{u\in\mathbb{F}_p^*}\sum_{x\in\mathbb{F}_{p^m}} \left(\zeta_p^{u\mathrm{Tr}_1^m((a+b)x^{p^k+1})} + \zeta_p^{u\mathrm{Tr}_1^m((a-b)\pi^{\frac{p^k+1}{2}}x^{p^k+1})}\right)\\ &= \sum_{u\in\mathbb{F}_p^*}\sum_{x\in\mathbb{F}_{p^m}} \left(\zeta_p^{u\mathrm{Tr}_1^m((a+b)x^2)} + \zeta_p^{u\mathrm{Tr}_1^m((a-b)\pi^{\frac{p^k+1}{2}}x^2)}\right)\end{aligned}$$

$$= \begin{cases} \sum\limits_{u\in\mathbb{F}_p^*}\sum\limits_{x\in\mathbb{F}_{p^m}}\left(\zeta_p^{u\mathrm{Tr}_1^m((a+b)x^2)}+\zeta_p^{u\mathrm{Tr}_1^m((a-b)x^2)}\right), & \text{if } p^k\equiv 3 \pmod 4, \\ \sum\limits_{u\in\mathbb{F}_p^*}\sum\limits_{x\in\mathbb{F}_{p^m}}\left(\zeta_p^{u\mathrm{Tr}_1^m((a+b)x^2)}+\zeta_p^{u\mathrm{Tr}_1^m((a-b)\pi x^2)}\right), & \text{if } p^k\equiv 1 \pmod 4. \end{cases}$$

If $p^k \equiv 1 \pmod 4$, then $\Delta_1 = \Delta_{\frac{p^k+1}{2}}$. On the other hand, if $p^k \equiv 3 \pmod 4$, then k is odd. Since s is odd, then m is odd, which implies that u_p is a nonsquare element in $\mathbb{F}_{p^m}$. In the following, we only need to prove that

$$\sum_{u\in\mathbb{F}_p^*}\sum_{x\in\mathbb{F}_{p^m}}\zeta_p^{u\mathrm{Tr}_1^m((a-b)x^2)} = \sum_{u\in\mathbb{F}_p^*}\sum_{x\in\mathbb{F}_{p^m}}\zeta_p^{u\mathrm{Tr}_1^m((a-b)\pi x^2)}.$$

On the other hand, we have that

$$\begin{aligned} &\sum_{u\in\mathbb{F}_p^*}\sum_{x\in\mathbb{F}_{p^m}}\zeta_p^{u\mathrm{Tr}_1^m((a-b)x^2)} \\ =\ &\sum_{u\in SQ_p}\sum_{x\in\mathbb{F}_{p^m}}\left(\zeta_p^{\mathrm{Tr}_1^m((a-b)(u^{\frac{1}{2}}x)^2)}+\zeta_p^{\mathrm{Tr}_1^m(u_p(a-b)(u^{\frac{1}{2}}x)^2)}\right) \\ =\ &\frac{p-1}{2}\sum_{x\in\mathbb{F}_{p^m}}\left(\zeta_p^{\mathrm{Tr}_1^m((a-b)x^2)}+\zeta_p^{\mathrm{Tr}_1^m(u_p(a-b)x^2)}\right) \\ =\ &\frac{p-1}{2}\sum_{x\in\mathbb{F}_{p^m}}\left(\zeta_p^{\mathrm{Tr}_1^m((a-b)x^2)}+\zeta_p^{\mathrm{Tr}_1^m(\pi^{\frac{p^m-1}{p-1}}(a-b)x^2)}\right) \\ =\ &\frac{p-1}{2}\sum_{x\in\mathbb{F}_{p^m}}\left(\zeta_p^{\mathrm{Tr}_1^m((a-b)x^2)}+\zeta_p^{\mathrm{Tr}_1^m((a-b)\pi(\pi^{\frac{p^{m-1}+p^{m-2}+\cdots+p}{2}}x)^2)}\right) \\ =\ &\frac{p-1}{2}\sum_{x\in\mathbb{F}_{p^m}}\left(\zeta_p^{\mathrm{Tr}_1^m((a-b)x^2)}+\zeta_p^{\mathrm{Tr}_1^m((a-b)\pi x^2)}\right), \end{aligned} \tag{3.8}$$

and

$$\begin{aligned} &\sum_{u\in\mathbb{F}_p^*}\sum_{x\in\mathbb{F}_{p^m}}\zeta_p^{u\mathrm{Tr}_1^m((a-b)\pi x^2)} \\ =\ &\sum_{u\in SQ_p}\sum_{x\in\mathbb{F}_{p^m}}\left(\zeta_p^{\mathrm{Tr}_1^m((a-b)\pi(u^{\frac{1}{2}}x)^2)}+\zeta_p^{\mathrm{Tr}_1^m(u_p(a-b)\pi(u^{\frac{1}{2}}x)^2)}\right) \\ =\ &\frac{p-1}{2}\sum_{x\in\mathbb{F}_{p^m}}\left(\zeta_p^{\mathrm{Tr}_1^m((a-b)\pi x^2)}+\zeta_p^{\mathrm{Tr}_1^m(u_p(a-b)\pi x^2)}\right) \\ =\ &\frac{p-1}{2}\sum_{x\in\mathbb{F}_{p^m}}\left(\zeta_p^{\mathrm{Tr}_1^m((a-b)\pi x^2)}+\zeta_p^{\mathrm{Tr}_1^m(\pi^{\frac{p^m-1}{p-1}+1}(a-b)x^2)}\right) \\ =\ &\frac{p-1}{2}\sum_{x\in\mathbb{F}_{p^m}}\left(\zeta_p^{\mathrm{Tr}_1^m((a-b)\pi x^2)}+\zeta_p^{\mathrm{Tr}_1^m((a-b)(\pi^{\frac{p^{m-1}+p^{m-2}+\cdots+p+2}{2}}x)^2)}\right) \end{aligned}$$

$$= \frac{p-1}{2} \sum_{x \in \mathbb{F}_{p^m}} \left(\zeta_p^{\mathrm{Tr}_1^m((a-b)\pi x^2)} + \zeta_p^{\mathrm{Tr}_1^m((a-b)x^2)} \right). \tag{3.9}$$

By comparing (3.8) and (3.9), we finish the proof. □

By Lemma 3.8, the weight distribution of the code $\mathcal{C}_{\frac{p^k+1}{2}}$ is the same as the code $\mathcal{C}_1$. As we know, the weight distribution of the code $\mathcal{C}_1$ has been studied in [14](see Theorems 5,6).

Lemma 3.9. [14] *With the notations given above.*

- *If $v_2(m) = 0$, then $\mathcal{C}_1$ is a cyclic code over $\mathbb{F}_p$ with parameters $[p^m - 1, 2m, \frac{p-1}{2}p^{m-1}]$ and the weight distribution of $\mathcal{C}_{\frac{p^k+1}{2}}$ is given in Table 3.*
- *If $1 \le v_2(m) \le v_2(k)$, then $\mathcal{C}_1$ is a cyclic code over $\mathbb{F}_p$ with parameters $[p^m - 1, 2m, \frac{p-1}{2}(p^{m-1} - p^{\frac{m}{2}-1})]$ and the weight distribution of $\mathcal{C}_{\frac{p^k+1}{2}}$ is given in Table 4.*

Table 3: for the case of $v_2(m) = 0$

Weight	Frequency
0	1
$\frac{p-1}{2}p^{m-1}$	$2(p^m - 1)$
$(p-1)p^{m-1}$	$p^{2m} - 2p^m + 1$

Table 4: for the case of $1 \le v_2(m) \le v_2(k)$

Weight	Frequency
0	1
$(p-1)p^{m-1}$	$\frac{(p^m-1)^2}{2}$
$(p-1)(p^{m-1} + p^{\frac{m}{2}-1})$	$\frac{(p^m-1)^2}{4}$
$(p-1)(p^{m-1} - p^{\frac{m}{2}-1})$	$\frac{(p^m-1)^2}{4}$
$\frac{p-1}{2}(p^{m-1} + p^{\frac{m}{2}-1})$	$p^m - 1$
$\frac{p-1}{2}(p^{m-1} - p^{\frac{m}{2}-1})$	$p^m - 1$

Applying Lemmas 3.8 and 3.9, we have the following theorem.

Theorem 3.10. *With the notations given above.*

- *If $v_2(m) = 0$, then $\mathcal{C}_{\frac{p^k+1}{2}}$ is a cyclic code over $\mathbb{F}_p$ with parameters $[p^m - 1, 2m, \frac{p-1}{2}p^{m-1}]$ and the weight distribution of $\mathcal{C}_{\frac{p^k+1}{2}}$ is given in Table 3.*
- *If $1 \le v_2(m) \le v_2(k)$, then $\mathcal{C}_{\frac{p^k+1}{2}}$ is a cyclic code over $\mathbb{F}_p$ with parameters $[p^m - 1, 2m, \frac{p-1}{2}(p^{m-1} - p^{\frac{m}{2}-1})]$ and the weight distribution of $\mathcal{C}_{\frac{p^k+1}{2}}$ is given in Table 4.*

3.3 The weight distribution of the code $\mathcal{C}_t$

Combining Lemma 3.1, Theorems 3.7 and 3.10, we have the following main result in this paper.

Theorem 3.11. *With the notations given above. Let $t \in \mathbb{Z}_{p^m-1}$ be a positive integer such that $t \equiv \frac{p^k+1}{2}p^\tau \pmod{\frac{p^m-1}{2}}$ for some $\tau \in \mathbb{Z}_m$, where k satisfies $\pi^{\frac{p^k+1}{2}p^i} \neq -\pi^{\frac{p^k+1}{2}}$ for all $i \in \mathbb{Z}_m$.*

- *If $v_2(k)+1 = v_2(m)$, then $\mathcal{C}_t$ is a cyclic code over $\mathbb{F}_p$ with parameters $[p^m-1, 2m, \frac{p-1}{2}(p^{m-1}-p^{\frac{m}{2}+d-1})]$ and the weight distribution of $\mathcal{C}_t$ is given by Table 1.*
- *If $v_2(k)+1 < v_2(m)$, then $\mathcal{C}_t$ is a cyclic code over $\mathbb{F}_p$ with parameters $[p^m-1, 2m, \frac{p-1}{2}(p^{m-1}-p^{\frac{m}{2}-1})]$ and the weight distribution of $\mathcal{C}_t$ is given by Table 2.*
- *If $v_2(m) = 0$, then $\mathcal{C}_t$ is a cyclic code over $\mathbb{F}_p$ with parameters $[p^m-1, 2m, \frac{p-1}{2}p^{m-1}]$ and the weight distribution of $\mathcal{C}_t$ is given by Table 3.*
- *If $1 \leq v_2(m) \leq v_2(k)$, then $\mathcal{C}_t$ is a cyclic code over $\mathbb{F}_p$ with parameters $[p^m-1, 2m, \frac{p-1}{2}(p^{m-1}-p^{\frac{m}{2}-1})]$ and the weight distribution of $\mathcal{C}_t$ is given by Table 4.*

In the following, we give two examples to verify our main results in Theorem 3.11.

Example 3.12. *Let $p = 3$, $m = 6$, $k = 1$. If $t \equiv 2p^\tau \pmod{364}$ for some $\tau \in \mathbb{Z}_6$, then the code $\mathcal{C}_t$ is a $[728, 12, 216]$ cyclic code over $\mathbb{F}_3$ with weight enumerator*

$$1 + 364X^{216} + 1092X^{252} + 33124X^{432} + 198744X^{468} + 298116X^{504},$$

which confirms the weight distribution in Table 1.

Example 3.13. *Let $p = 3$, $m = 4$, $k = 1$. If $t \equiv 2p^\tau \pmod{40}$ for some $\tau \in \mathbb{Z}_4$, then the code $\mathcal{C}_t$ is a $[80, 8, 24]$ cyclic code over $\mathbb{F}_3$ with weight enumerator*

$$1 + 120X^{24} + 40X^{36} + 3600X^{48} + 2400X^{60} + 400X^{72},$$

which confirms the weight distribution in Table 2.

Example 3.14. *Let $p = 5$, $m = 3$, $k = 1$. If $t \equiv 3p^\tau \pmod{62}$ for some $\tau \in \mathbb{Z}_3$, then the code $\mathcal{C}_t$ is a $[124, 6, 50]$ cyclic code over $\mathbb{F}_5$ with weight enumerator*

$$1 + 248X^{50} + 15376X^{100},$$

which confirms the weight distribution in Table 3.

Example 3.15. *Let $p = 3$, $m = 6$, $k = 2$. If $t \equiv 5p^\tau \pmod{364}$ for some $\tau \in \mathbb{Z}_6$, then the code $\mathcal{C}_t$ is a $[728, 12, 234]$ cyclic code over $\mathbb{F}_3$ with weight enumerator*

$$1 + 728X^{234} + 728X^{252} + 132496X^{468} + 264992X^{486} + 132496X^{504},$$

which confirms the weight distribution in Table 4.

References

[1] P. Delsarte, "On subfield subcodes of modified Reed-Solomon codes," *IEEE Trans. Inform. Theory*, **21**(5), 575-576(1975).

[2] C. Ding, Y. Liu, C. Ma, L. Zeng, "The weight distributions of the duals of cyclic codes with two zeros," *IEEE Trans. Inform. Theory*, **57**(12), 8000-8006(2011).

[3] C. Ding, J. Yang, "Hamming weight in irrecducible codes," *Discret. Math.* **313**(4), 434-446(2013).

[4] S. Draper, X. Hou, "explicit evaluation of certain exponential sums of quadratic functions over $\mathbb{F}_{p^m}$, p odd," http://arxiv.org/pdf/0708.3619v1.pdf.

[5] K. Feng, J. Luo, "Weight distribution of some reducible cyclic codes," *Finite Fields Appl.* **14**(2), 390-409(2008).

[6] T. Feng, "On cyclic codes of length $2^{2^r} - 1$ with two zeros whose dual codes have three weights," *Des. Codes Cryptogr.* **62**(3), 253-258(2012).

[7] T. Helleseth, A. Kholosha, "Monomial and quadratic bent functions over the finite fields of odd characteristic", *IEEE Trans. Inform. Theory,* **52**(5), 2018-2032(2006).

[8] A. Klapper, "Cross-correlations of quadratic form sequences in odd characteristic," *Des. Codes Cryptogr.* **3**(4), 289-305(1997).

[9] C. Li, N. Li, T. Helleseth, C. Ding, "On the weight distributions of several classes of cyclic codes from APN monomials,"Available: http://arxiv.org/pdf/1308.5885v1.pdf.

[10] R. Lidl, H. Niederreiter, "Finite Fields," Ency clopedia of Mathematics, **20**, Cambridge University Press, Cambridge(1983).

[11] Y. Liu, H. Yan, C. Liu, "A class of six-weight cyclic codes and their weight distribution," Available: http://arxiv.org/pdf/1311.3391v2.pdf.

[12] J. Luo, K. Feng, "Cyclic codes and sequences from generalized Coulter-Matthews function," *IEEE Trans. Inform. Theory*, **54**(12), 5345-5353(2008).

[13] J. Luo, K. Feng, "On the weight distributions of two classes of cyclic codes," *IEEE Trans. Inform. Theory*, **54**(12), 5332-5344(2008).

[14] C. Ma, L. zeng, Y. Liu, D. Feng, C. Ding, "The weight enumerator of a class of cyclic codes," *IEEE Trans. Inform. Theory*, **57**(1), 397-402(2011).

[15] F. MacWilliams, N. Sloane, "The Theory of Error-Correcting Codes," North-Holland Publishing, Amsterdam(1997).

[16] A. Rao, N. Pinnawala, "A family of two-weight irreducible cyclic codes," *IEEE Trans. Inform. Theory*, **56**(6), 2568-2570(2010).

[17] H. Trachtenberg, "On the crosscorrelation functions of maximal linear recurring sequences," Ph.D. dissertation, Univ. South. Calif., Los Angels(1970).

[18] G. Vega, "The weight distribution of an extended class of reducible cyclic codes," *IEEE Trans. Inform. Theory*, **58**(7), 4862-4869(2012).

[19] G. Vega, J. Wolfmann, "New classes of 2-weight cyclic codes," *Des. Codes Cryptogr.* **42**(3), 327-334(2007).

[20] B. Wang, C. Tang, Y. Qi , Y. Yang , M. Xu, "The weight distributions of cyclic codes and elliptic curves," *IEEE Trans. Inform. Theory*, **58**(12), 7253-7259(2012).

[21] M. Xiong, "The weight distributions of a class of cyclic codes II," *Des. Codes Cryptogr.* (2012). doi: 10.1007/s10623-012-9785-0.

[22] M. Xiong, "The weight distributions of a class of cyclic codes III," *Finite Fields Appl.* **21**, 84-96(2013).

[23] M. Xiong, "The weight distributions of a class of cyclic codes," *Finite Fields Appl.* **18**(5), 933-945(2012).

[24] X. Zeng, J. Shan, L. Hu, "A triple-error-correcting cyclic code from the Gold and Kasami-Welch APN power functions," *Finite Fields Appl.* **18**(1), 70-92(2012).

[25] X. Zeng, L. Hu, W. Jiang, Q. Yue, X. Cao, "The weight distribution of a class of p-ary cyclic codes," *Finite Fields Appl.* **16**(1), 56-73(2010).

[26] D. Zheng, X. Wang, X. Zeng, L. Hu, "The weight distribution of a family of p-ary cyclic codes," Des. Codes Cryptogr. doi: 10.1007/s10623-013-9908-2.

[27] Z. Zhou, C. Ding, "A class of three-weight cyclic codes," *Finite Fields Appl.* **25**, 79-93(2014).

[28] Z. Zhou, C. Ding, J. Luo, A. Zhang, "A family of five-weight cyclic codes and their weight enumerators," *IEEE Trans. Inform. Theory*, **59**(10), 6674-6682(2013).